arXiv:cond-mat/0404567v2 [cond-mat.soft] 1 Jun 2004

Berry phase caused by nondissipative drag of superflow in a Bose qubit

D V Fil†§ and S I Shevchenko‡

†Institute for Single Crystals National Academy of Sciences of Ukraine, Lenin av. 60, Kharkov 61001, Ukraine
‡Verkin Institute for Low Temperature Physics and Engineering National Academy of Sciences of Ukraine, Lenin av. 47, Kharkov 61103, Ukraine

Abstract. A microscopic theory of superfluid drag in a two-component Bose gas is developed. The drag factor is shown to be proportional to the square root of the gas parameter. Basing on the similarity between the "drag force" and vector potential of magnetic field we propose how the superfluid drag can be used to detect a Berry phase in a Bose counterpart of the Josephson charge qubit. We consider a system in which the drag component, situated inside the drive component, is confined in two half-ring traps separated by two Josephson barriers. Under cyclic adiabatic change of Josephson coupling parameters and an asymmetry of two traps a Berry phase is generated. This phase can be observed though measurements of relative number of atoms in two traps. The Berry phase depends on the drag factor and its detection can be used for determining the value of the drag.

PACS numbers: 03.75.Kk, 03.75.Lm, 03.67.Lx

---

§ fil@isc.kharkov.com

1. Introduction

The interest in studying of macroscopic quantum phenomena in Josephson superconductive devices has renewed recently in connection with the possibility of using such devices for performing quantum logic operations [1]. Related problem - Josephson-type effects in Bose systems was received considerable attention [2] after experimental realization of Bose-Einstein condensation in ultracold atomic gases. It was shown in [3] that Josephson Bose system can also be used for quantum computing. In [3] a simple model of a Bose counterpart of the Josephson charge qubit has been formulated.

The supercurrent in superconductors is coupled to the vector potential of electromagnetic fields that allows to perform quantum logic operations by external magnetic fields. External magnetic fields couple with Bose atoms only through Zeeman and hyperfine interactions. Therefore, in difference with the case of superconductors, magnetic field cannot be used to control supercurrents in uncharged Bose systems.

In this paper we consider the effect that can be used for quantum state engineering in Bose Josephson systems in the same manner as magnetic field in superconductive Josephson qubits. This effect is a nondissipative drag of supercurrent in a two-component superfluid system.

Nondissipative drag is caused by interaction between the atoms and consists in a redistribution of a supercurrent between two superfluid components. For the first time the nondissipative drag was considered by Andreev and Bashkin [4]. In [4] a three-velocity hydrodynamic model of a ^{3}He-^{4}He mixture was developed. It was shown that superfluid behavior of such a system can be described under accounting the "drag" term in the free energy. This term is proportional to the scalar product of the superfluid velocities of two components times the difference between the effective and the bare masses of ^{3}He atoms. A microscopic theory of the nondissipative drag for a special case of a bilayer system of charged bosons was developed in Refs. [5, 6].

Nondissitive drag is a fundamental property of systems with a macroscopic quantum coherence and takes place also in superconductors [7], quantum Hall systems [8] and quantum wires [9]. But for the system of uncharged bosons this effect is especially important since the "drag force" plays the role similar to the role of the vector potential of a magnetic field in superconductors [4].

An important problem of quantum computation is to minimize the influence of random noise coming from the environment. One of the ways of solutions of this problem consists in using of geometrical phases to implement quantum gates [10]. As was shown in [11] a geometrical Berry phase can be generated in a SQUID loop with two Josephson junctions in an external magnetic field. In this paper we propose a model of Bose qubit that may demonstrate analogous effect.

In Sec. 2 the theory of nondissipative drag in two-component Bose gases is developed. In Sec. 3 the model of Bose qubit is formulated. The scheme of generating of the Berry phase is proposed in Sec. 4. Conclusions are given in Sec. 5.

2. Nondissipative drag of superflow in a two-component Bose system

In this section we develop a microscopic theory of the nondissipative drag in a two-component atomic Bose system with point two-particle interaction.

Let us consider an infinite uniform two-component atomic Bose gas in a Bose-Einstein condensed state. In general case the densities of atoms in each component are different from one another ($n_1 \neq n_2$), the atoms of each components have different masses ($m_1 \neq m_2$) and the interaction between atoms is described by three different scattering lengths ($a_{11} \neq a_{22} \neq a_{12}$). The Hamiltonian of the system can be presented in the form

$$H = \sum_{i=1,2} (E_i - \mu_i N_i) + \frac{1}{2} \sum_{i,i'=1,2} E_{ii'}^{int}, \tag{1}$$

where

$$E_i = \int d^3r \frac{\hbar^2}{2m_i} [\nabla \hat{\Psi}_i^+(\mathbf{r})] \nabla \hat{\Psi}_i(\mathbf{r}) \tag{2}$$

is the kinetic energy,

$$E_{ii'}^{int} = \int d^3r \hat{\Psi}_i^+(\mathbf{r}) \hat{\Psi}_{i'}^+(\mathbf{r}) \gamma_{ii'} \hat{\Psi}_{i'}(\mathbf{r}) \hat{\Psi}_i(\mathbf{r}) \tag{3}$$

is the energy of interaction, $\gamma_{ii} = 4\pi\hbar^2 a_{ii}/m_i$ and $\gamma_{12} = 2\pi\hbar^2(m_1 + m_2)a_{12}/(m_1 m_2)$ are the interaction parameters.

For further analysis it is convenient to use the density and phase operator approach (see, for instance, [12, 13]). The approach is based on the following representation for the Bose field operators

$$\hat{\Psi}_i(\mathbf{r}) = \exp\left[i\varphi_i(\mathbf{r}) + i\hat{\varphi}_i(\mathbf{r})\right] \sqrt{n_i + \hat{n}_i(\mathbf{r})}, \tag{4}$$

$$\hat{\Psi}_i^+(\mathbf{r}) = \sqrt{n_i + \hat{n}_i(\mathbf{r})} \exp\left[-i\varphi_i(\mathbf{r}) - i\hat{\varphi}_i(\mathbf{r})\right], \tag{5}$$

where $\hat{n}_i$ and $\hat{\varphi}_i$ are the density and phase fluctuation operators, $\varphi_i(\mathbf{r})$ is the c-number term of the phase operator, which are connected with the superfluid velocity by common relation $\mathbf{v}_i = \hbar\nabla\varphi_i/m_i$. In what follows we specify the case of superfluid velocities independent of $\mathbf{r}$.

Substituting Eqs. (4), (5) into Eq. (1) and expanding it in series in powers of $\hat{n}_i$ and $\nabla\hat{\varphi}_i$ we present the Hamiltonian of the system in the following form

$$H = H_0 + H_2 + \ldots \tag{6}$$

In (6) the term

$$H_0 = V\left(\sum_{i=1,2}\left[\frac{1}{2}m_i n_i \mathbf{v}_i^2 + \frac{\gamma_{ii}}{2}n_i^2 - \mu_i n_i\right] + \gamma_{12} n_1 n_2\right) \tag{7}$$

does not contain the operator part (V is the volume of the system). The chemical potentials are determined from the minimization condition of the Hamiltonian H_0:

$$\frac{1}{2}m_i \mathbf{v}_i^2 + \gamma_{ii} n_i + \gamma_{12} n_{3-i} - \mu_i = 0. \tag{8}$$

Fulfillment of Eq. (8) results in vanishing of linear in the density fluctuation operators terms in the Hamiltonian. The terms linear in the phase fluctuation operators are absent in the Hamiltonian due to continuity conditions.

The quadratic in $\nabla\hat{\varphi}_i$ and $\hat{n}_i$ operator term in H reads as

$$H_2 = \int d\mathbf{r}\Bigg(\sum_i \Bigg\{\frac{\hbar^2}{2m_i}\left[\frac{\Big(\nabla\hat{n}_i(\mathbf{r})\Big)^2}{4n_i} + n_i\Big(\nabla\hat{\varphi}_i(\mathbf{r})\Big)^2\right]$$
$$+\frac{\hbar\mathbf{v}_i}{2}\Big(\hat{n}_i(\mathbf{r})\nabla\hat{\varphi}_i(\mathbf{r}) + [\nabla\hat{\varphi}_i(\mathbf{r})]\hat{n}_i(\mathbf{r})\Big)$$

$$+\frac{i\hbar^2}{2m_i}\Big([\nabla\hat{n}_i(\mathbf{r})]\nabla\hat{\varphi}_i(\mathbf{r})-[\nabla\hat{\varphi}_i(\mathbf{r})]\nabla\hat{n}_i(\mathbf{r})\Big)+\frac{\gamma_{ii}}{2}\Big(\hat{n}_i(\mathbf{r})\Big)^2\Bigg\}+\gamma_{12}\hat{n}_1(\mathbf{r})\hat{n}_2(\mathbf{r})\Bigg). \qquad (9)$$

The quadratic part of the Hamiltonian determines the spectrum of the elementary excitations. Hereafter we will neglect the higher order terms in the Hamiltonian (6). These terms describe the scattering of the quasiparticles and they can be omitted if the temperature is much smaller than the temperature of Bose-Einstein condensation.

Let us rewrite the quadratic part of the Hamiltonian in terms of the operators of creation and annihilation of the elementary excitations. As the first step we use the substitution

$$\hat{n}_i(\mathbf{r})=\sqrt{\frac{n_i}{V}}\sum_{\mathbf{k}}e^{i\mathbf{kr}}\sqrt{\frac{\epsilon_{ik}}{E_{ik}}}\left[b_i(\mathbf{k})+b_i^+(-\mathbf{k})\right], \qquad (10)$$

$$\hat{\varphi}_i(\mathbf{r})=\frac{1}{2i}\sqrt{\frac{1}{n_iV}}\sum_{\mathbf{k}}e^{i\mathbf{kr}}\sqrt{\frac{E_{ik}}{\epsilon_{ik}}}\left[b_i(\mathbf{k})-b_i^+(-\mathbf{k})\right], \qquad (11)$$

where operators b_i^+, b_i satisfy the Bose commutative relations. Here $\epsilon_{ik}=\hbar^2k^2/2m_i$ is the spectrum of free atoms, and $E_{lk}=\sqrt{\epsilon_{ik}(\epsilon_{ik}+2\gamma_{ii}n_i)}$ is the spectrum of the elementary excitations at $\gamma_{12}=0$ and $\mathbf{v}_i=0$. The substitution (10), (11) reduces the Hamiltonian (9) to the form, quadratic in operators b_i^+, b_i:

$$H_2=\sum_{i\mathbf{k}}\left[\mathcal{E}_i(\mathbf{k})\left(b_i^+(\mathbf{k})b_i(\mathbf{k})+\frac{1}{2}\right)-\frac{1}{2}\epsilon_{ik}\right]$$
$$+\sum_{\mathbf{k}}g_k\left[b_1^+(\mathbf{k})b_2(\mathbf{k})+b_1(\mathbf{k})b_2(-\mathbf{k})+h.c.\right]. \qquad (12)$$

Here

$$\mathcal{E}_i(\mathbf{k})=E_{ik}+\hbar\mathbf{k}\mathbf{v}_i \qquad (13)$$

and

$$g_k=\gamma_{12}\sqrt{\frac{\epsilon_{1k}\epsilon_{2k}n_1n_2}{E_{1k}E_{2k}}}. \qquad (14)$$

The Hamiltonian (12) contains non-diagonal in Bose creation and annihilation operator terms and it can be diagonalized using the standard procedure of u-v transformation [14]. The result is

$$H_2=\sum_{\mathbf{k}}\left[\sum_{\lambda=\alpha,\beta}\mathcal{E}_\lambda(\mathbf{k})\left(\beta_\lambda^+(\mathbf{k})\beta_\lambda(\mathbf{k})+\frac{1}{2}\right)-\frac{1}{2}\sum_{i=1,2}\epsilon_{ik}\right], \qquad (15)$$

where $\beta_\lambda(k)^+$ and $\beta_\lambda(k)$ are the operators of creation and annihilation of elementary excitations.

The energies $\mathcal{E}_\lambda(\mathbf{k})$ are found as solutions of the equation

$$\det\begin{pmatrix}\mathbf{A}-\mathcal{E}\mathbf{I} & \mathbf{B}\\ \mathbf{B} & \mathbf{A}+\mathcal{E}\mathbf{I}\end{pmatrix}=0, \qquad (16)$$

where

$$\mathbf{A}=\begin{pmatrix}\mathcal{E}_1(\mathbf{k}) & 0 & g_k & 0\\ 0 & \mathcal{E}_1(-\mathbf{k}) & 0 & g_k\\ g_k & 0 & \mathcal{E}_2(\mathbf{k}) & 0\\ 0 & g_k & 0 & \mathcal{E}_2(-\mathbf{k})\end{pmatrix} \qquad (17)$$

$$\mathbf{B}=\begin{pmatrix} 0 & 0 & 0 & g_k \\ 0 & 0 & g_k & 0 \\ 0 & g_k & 0 & 0 \\ g_k & 0 & 0 & 0 \end{pmatrix} \tag{18}$$

and $\mathbf{I}$ is the identity matrix. Having the spectra of elementary excitations one can obtain the relation between the supercurrents in each component are their superfluid velocities.

We will find the current densities using the relation

$$\mathbf{j}_i=\frac{1}{V}\frac{\partial F}{\partial \mathbf{v}_i}, \tag{19}$$

where F is the free energy of the system. Here the quantity $\mathbf{j}_i$ is defined as density of the mass current.

The free energy of the system, described by the Hamiltonian (6), is given by the formula

$$F=H_0+\frac{1}{2}\sum_{\mathbf{k}}\left[\sum_{\lambda=\alpha,\beta}\mathcal{E}_\lambda(\mathbf{k})-\sum_{i=1,2}\epsilon_{ik}\right]+T\sum_{\mathbf{k}}\sum_{\lambda=\alpha,\beta}\ln\left[1-\exp\left(-\frac{\mathcal{E}_\lambda(\mathbf{k})}{T}\right)\right]. \tag{20}$$

The second term in (20) is the energy of the zero-point fluctuations and the third term is the standard temperature dependent part of the free energy for the gas of noninteracting elementary excitations.

To proceed further we consider the case of small superfluid velocities, much smaller than the velocities of elementary excitations, and seek for the free energy as series in $\mathbf{v}_i$.

At $\mathbf{v}_1=\mathbf{v}_2=0$ the equation (16) is easily solved and the spectra are found to be

$$E_{\alpha(\beta)k}=\left(\frac{E_{1k}^2+E_{2k}^2}{2}\pm\sqrt{\frac{(E_{1k}^2-E_{2k}^2)^2}{4}+4\gamma_{12}^2 n_1 n_2\epsilon_{1k}\epsilon_{2k}}\right)^{1/2}. \tag{21}$$

As required in the procedure [14] we take positive valued solutions of Eq. (16). We should emphasize that the energies (21) should be real valued. This requirement yields the common condition of stability of a two-component system: $\gamma_{12}^2\leq\gamma_{11}\gamma_{22}$. If this condition were not fulfilled spatial separation of two components (at positive γ_{12}) or a collapse (at negative γ_{12}) would take place.

At nonzero superfluid velocities we present the solutions of Eq. (16) as series in $\mathbf{v}_i$, neglecting the terms higher than quadratic:

$$\mathcal{E}_\alpha(\mathbf{k})=E_{\alpha k}+\frac{1}{2}\hbar\mathbf{k}\mathbf{v}_1\left(1+\frac{E_{1k}^2-E_{2k}^2}{E_{\alpha k}^2-E_{\beta k}^2}\right)+\frac{1}{2}\hbar\mathbf{k}\mathbf{v}_2\left(1-\frac{E_{1k}^2-E_{2k}^2}{E_{\alpha k}^2-E_{\beta k}^2}\right)$$
$$+\frac{2\gamma_{12}^2 n_1 n_2\epsilon_{1k}\epsilon_{2k}\left(3E_{\alpha k}^2+E_{\beta k}^2\right)}{E_{\alpha k}\left(E_{\alpha k}^2-E_{\beta k}^2\right)^3}\hbar^2\left(\mathbf{k}\mathbf{v}_1-\mathbf{k}\mathbf{v}_2\right)^2, \tag{22}$$

$$\mathcal{E}_\beta(\mathbf{k})=E_{\beta k}+\frac{1}{2}\hbar\mathbf{k}\mathbf{v}_1\left(1-\frac{E_{1k}^2-E_{2k}^2}{E_{\alpha k}^2-E_{\beta k}^2}\right)+\frac{1}{2}\hbar\mathbf{k}\mathbf{v}_2\left(1+\frac{E_{1k}^2-E_{2k}^2}{E_{\alpha k}^2-E_{\beta k}^2}\right)$$
$$-\frac{2\gamma_{12}^2 n_1 n_2\epsilon_{1k}\epsilon_{2k}\left(E_{\alpha k}^2+3E_{\beta k}^2\right)}{E_{\alpha k}\left(E_{\alpha k}^2-E_{\beta k}^2\right)^3}\hbar^2\left(\mathbf{k}\mathbf{v}_1-\mathbf{k}\mathbf{v}_2\right)^2. \tag{23}$$

Note that at $\mathbf{v}_1 = \mathbf{v}_2 = \mathbf{v}$ the spectra (22), (23) are reduced to common expressions for the energies of quasiparticles in a moving condensate: $\mathcal{E}_{\alpha(\beta)}(\mathbf{k}) = E_{\alpha(\beta)k} + \hbar\mathbf{k}\mathbf{v}$.

Using Eqs. (20),(22) and (23) we obtain the following expression for the free energy

$$F = F_0 + \frac{V}{2}\left[(\rho_1 - \rho_{n1})\mathbf{v}_1^2 + (\rho_2 - \rho_{n2})\mathbf{v}_2^2 - \rho_{\rm dr}(\mathbf{v}_1 - \mathbf{v}_2)^2\right], \qquad (24)$$

where F_0 does not depend on $\mathbf{v}_i$. In (24) $\rho_i = m_i n_i$ are the mass densities, the quantities

$$\rho_{n1} = -\frac{2m_1}{3V}\sum_{\mathbf{k}}\epsilon_{1k}\left[\frac{dN_{\alpha k}}{dE_{\alpha k}} + \frac{dN_{\beta k}}{dE_{\beta k}} + \frac{E_{1k}^2 - E_{2k}^2}{E_{\alpha k}^2 - E_{\beta k}^2}\left(\frac{dN_{\alpha k}}{dE_{\alpha k}} - \frac{dN_{\beta k}}{dE_{\beta k}}\right)\right], \qquad (25)$$

$$\rho_{n2} = -\frac{2m_2}{3V}\sum_{\mathbf{k}}\epsilon_{2k}\left[\frac{dN_{\alpha k}}{dE_{\alpha k}} + \frac{dN_{\beta k}}{dE_{\beta k}} - \frac{E_{1k}^2 - E_{2k}^2}{E_{\alpha k}^2 - E_{\beta k}^2}\left(\frac{dN_{\alpha k}}{dE_{\alpha k}} - \frac{dN_{\beta k}}{dE_{\beta k}}\right)\right] \qquad (26)$$

describe the thermal reduction of the superfluid densities, and the quantity

$$\rho_{dr} = \frac{4}{3V}\sqrt{m_1 m_2}\sum_{\mathbf{k}}\frac{\gamma_{12}^2 n_1 n_2\left(\epsilon_{1k}\epsilon_{2k}\right)^{3/2}}{E_{\alpha k}E_{\beta k}}\left[\frac{1 + N_{\alpha k} + N_{\beta k}}{(E_{\alpha k} + E_{\beta k})^3} - \frac{N_{\alpha k} - N_{\beta k}}{(E_{\alpha k} - E_{\beta k})^3}\right.$$
$$\left. + \frac{2E_{\alpha k}E_{\beta k}}{\left(E_{\alpha k}^2 - E_{\beta k}^2\right)^2}\left(\frac{dN_{\alpha k}}{dE_{\alpha k}} + \frac{dN_{\beta k}}{dE_{\beta k}}\right)\right], \qquad (27)$$

which we call the "drag density," yields the value of redistribution of the superfluid densities between components. In Eqs. (25)-(27) $N_{\alpha(\beta)k} = [\exp(E_{\alpha(\beta)k}/T) - 1]^{-1}$ is the Bose distribution function.

Using Eqs. (19), (24) we arrive to the following expressions for the supercurrents

$$\mathbf{j}_1 = (\rho_1 - \rho_{n1} - \rho_{dr})\mathbf{v}_1 + \rho_{\rm dr}\mathbf{v}_2, \qquad (28)$$

$$\mathbf{j}_2 = (\rho_2 - \rho_{n2} - \rho_{dr})\mathbf{v}_2 + \rho_{\rm dr}\mathbf{v}_1. \qquad (29)$$

One can see that at nonzero ρ_{dr} the current of one component contains the term proportional to the superfluid velocity of the other component. It means that there is a transfer of motion between the components, or the drag effect. Since $\rho_{dr} \propto \gamma_{12}^2$ the effect occurs only at nonzero interaction between the components and it does not depend on the sign of this interaction.

Eq. (27) is the main result of this section. This equation yields the value of the drag for a general case of two-component Bose system with components having different densities, different masses of atoms, different interaction parameters, and for zero as well as for nonzero temperatures.

To estimate the absolute value of the drag we, for simplicity, specify the case of $m_1 = m_2$, that is realized when two components are two hyperfine states of the same atoms.

At $m_1 = m_2 = m$ and $T = 0$ Eq. (27) is reduced to the form

$$\rho_{dr} = \frac{4m}{3}\int_0^\infty d\epsilon\frac{\gamma_{12}^2 n_1 n_2 \nu(\epsilon)\epsilon^{1/2}}{\sqrt{(\epsilon + w_1)(\epsilon + w_2)}\left(\sqrt{\epsilon + w_1} + \sqrt{\epsilon + w_2}\right)^3}, \qquad (30)$$

where

$$w_{1(2)} = \gamma_{11}n_1 + \gamma_{22}n_2 \pm \sqrt{(\gamma_{11}n_1 - \gamma_{22}n_2)^2 + 4\gamma_{12}^2 n_1 n_2} \qquad (31)$$

and

$$\nu(\epsilon) = \frac{m^{3/2}}{\sqrt{2}\pi^2\hbar^3}\sqrt{\epsilon}$$

is the density of states of elementary excitations.

The integral in (30) is evaluated analytically. To present the answer in a compact form it is convenient to introduce dimensionless parameters

$$\eta = \frac{\gamma_{12}^2}{\gamma_{11}\gamma_{22}} \quad \text{and} \quad \kappa = \sqrt{\frac{n_1\gamma_{11}}{n_2\gamma_{22}}}.$$

Using these notations we have

$$\rho_{dr} = \sqrt{\rho_1\rho_2}\sqrt[4]{n_1 a_{11}^3 n_2 a_{22}^3}\,\eta^2 F(\kappa,\eta), \tag{32}$$

where

$$F(\kappa,\eta) = \frac{256}{45\sqrt{2\pi}}\left(\kappa+\kappa^{-1}+3\sqrt{1-\eta^2}\right) \times\left(\sqrt{\kappa+\kappa^{-1}+\sqrt{(\kappa-\kappa^{-1})^2+4\eta^2}}+\sqrt{\kappa+\kappa^{-1}-\sqrt{(\kappa-\kappa^{-1})^2+4\eta^2}}\right)^{-3}. \tag{33}$$

At $a_{11}n_1 = a_{22}n_2$ we obtain from (32) the following approximate relation

$$\frac{\rho_{dr}}{\rho_1} \approx \frac{1}{2}\frac{a_{12}^2}{a_{11}a_{22}}\sqrt{n_1 a_{11}^3}. \tag{34}$$

For $n_2 \gg n_1$ and $a_{11} \sim a_{22}$ we find from (32) that

$$\frac{\rho_{dr}}{\rho_1} \approx 0.8\frac{a_{12}^2}{a_{22}^2}\sqrt{n_2 a_{22}^3}. \tag{35}$$

Since the scattering lengths in two-component systems are of the same order and the gas parameters $n_i a_{ii}^3$ can be as much as 10^{-4} the quantity ρ_{dr}/ρ_1 can be of order of 10^{-2}.

Eq. (27) shows that the "drag density" decreases under increasing the temperature. Evaluating the temperature dependence of ρ_{dr} numerically we conclude that at the temperatures of order of the interaction energy this decreasing is not crucial. As an example, the dependence of ρ_{dr} on the temperature at $n_1 = n_2 = n$ and $\gamma_{11} = \gamma_{22} = \gamma$ is shown in Fig. 1. Results, presented in Fig. 1, correspond to $\eta = 0.5$, but, actually, temperature dependence of ρ_{dr} is not very sensitive to this parameter.

In conclusion, let us discuss to which extend the results, obtained in this section, can be applied to Bose gases confined in a trap. In nonuniform systems the spectrum of elementary excitations is modified [13, 15, 16, 17], but for healing lengths much smaller than the linear size of the Bose clouds the spectrum of excitations at wave vectors of order or higher than inverse healing length is well described by the quasi uniform approximation. The main contribution to the sum in Eq. (27) comes from the excitations with such wave vectors. Therefore, the drag effect in nonuniform system can be described by the same equations, as in the uniform case with the only modification that the quantities n_1 and n_2, and, correspondingly, ρ_i, ρ_{ni}, ρ_{dr} and j_i in Eqs. (25)-(29) are understood as functions of coordinates.



Figure 1. Dependence of the "drag density" on the temperature.

3. Model of BEC qubit with external drag force

Let our two-component system is confined in a toroidal trap and the Bose cloud of component 1 (drag component) is situated inside and overlaps with the Bose clouds of component 2 (drive component). Such a situation can be realized if $|\gamma_{12}| < \min(\gamma_{11}, \gamma_{22})$. Then the confining potential is deformed to cut the drag component in two clouds of a half-torus shape (separated by two Josephson links), but the toroidal shape of the Bose cloud of the drive component is survived under such a deformation (see Fig. 2). In what follows we imply that circumference $L = 2\pi R_t$ of the toroidal clouds is much larger than radiuses of their cross-sections r_{t1} and r_{t2}.

Rotating this trap one can excite a superflow in the drive component. After the rotation be switched off there will be a circulating superflow in the drive component and no superflow in the drag component (at negligible small Josephson coupling). The superfluid velocity of the drive component satisfies the Feynman-Onsager quantization conditions and its value is proportional to the vorticity parameter N_v for the drive component:

$$v_2 = \frac{N_v \hbar}{m_2 R_t} \tag{36}$$

(N_v is integer). In (36) we, for simplicity, neglect the effect caused by variation of r_{t2}.

Since $j_1 = 0$, the phase gradient $\nabla\varphi_1$ should be nonzero to compensate the drag effect. In polar coordinates the ϕ component of the phase gradient is given by the



Figure 2. Schematic shapes of Bose clouds for the drag (top figure) and drive (bottom figure) components. In system considered the drag component overlaps with the drive component.

relation

$$(\nabla\varphi_1)_\phi = -\frac{N_v}{R_t} f_{dr}, \tag{37}$$

where

$$f_{dr} = \frac{m_1}{m_2}\frac{\rho_{dr}}{\rho_1 - \rho_{n1} - \rho_{dr}} \tag{38}$$

is the drag factor. We imply that r_{t1} and r_{t2} are much higher than healing lengths that allows to describe the drag effect in quasi-uniform approximation. For definiteness, we specify the case of $\rho_1 \ll \rho_2$ and $\rho_2 \approx const$ in the overlapping region. In such a case one can neglect space dependence the drag factor (see Eq. (35)).

At nonzero Josephson coupling the current j_1 can be nonzero, but it cannot exceed maximum Josephson current J_m. Relation (37) remains approximately correct at nonzero Josephson coupling, if a strong inequality $J_m \ll \hbar\rho_1/(mR_t)$ is satisfied. Here we specify just such a case.

The drag force can be considered as external "magnetic field" applied to the drag component. The latter component in the geometry considered forms a superfluid circuit with two Josephson links. The situation described is analogous to the one that takes place in superconductive Josephson qubits [1, 11] and our system can be treated as a Bose qubit controlled by the drag force.

We will formulate the model of "dragged" Bose qubit using an approach similar to one of Ref. [3].

Let us present the Bose field operators for the component 1 in the two mode approximation:

$$\begin{aligned} \hat{\Psi}_1(\mathbf{r},t) &= \sum_{l=L,R} a_l(t)\Psi_l(\mathbf{r}-\mathbf{r}_l), \\ \hat{\Psi}_1^+(\mathbf{r},t) &= \sum_{l=L,R} a_l^+(t)\Psi_l^*(\mathbf{r}-\mathbf{r}_l), \end{aligned} \tag{39}$$

where $a^+_{L(R)}$ and $a_{L(R)}$ are the operators of creation and annihilation of bosons in the condensate confined in the left(right) half-torus, and Ψ_L, Ψ_R are two almost orthogonal local mode functions

$$\int d^3r\Psi_l^*(\mathbf{r})\Psi_{l'}(\mathbf{r}) \approx \delta_{ll'}, \quad l,l' = L,R$$

that describe the condensate in the left and right traps [18].

Substituting (39) into Hamiltonian (1), we obtain the following expression for the parts of the Hamiltonian that depends on the operators a_l^+, a_l:

$$H_a = \sum_{l=L,R}\left(K_l a_l^+ a_l + \lambda_l a_l^+ a_l^+ a_l a_l\right) + (J a_L^+ a_R + J^* a_R^+ a_L). \tag{40}$$

Here

$$K_l = \int d^3r \Psi_l^*\left[-\frac{\hbar^2}{2m}\nabla^2 + V_{tr} + \gamma_{12}\Psi_2^*\Psi_2\right]\Psi_l, \tag{41}$$

$$\lambda_l = \frac{\gamma_{11}}{2}\int d^3r|\Psi_l|^4, \tag{42}$$

$$J = \int d^3r\left[\frac{\hbar^2}{2m}\nabla\Psi_L^*\nabla\Psi_R + V_{tr}\Psi_L^*\Psi_R\right]. \tag{43}$$

Functions Ψ_L and Ψ_R contain the phase factors $e^{i\varphi_L(\mathbf{r})}$ and $e^{i\varphi_R(\mathbf{r})}$ with the phases satisfying Eq. (37). Taking these factors into account, we choose the following basis for the one mode wave functions

$$\Psi_{L(R)}(\mathbf{r}) = |\Psi_{L(R)}(\mathbf{r})|\exp\left[i\phi_{L(R)}(\mathbf{r})\Phi_{dr}\right], \tag{44}$$

where

$$\Phi_{dr} = -N_v f_{dr} \tag{45}$$

and ϕ_L,ϕ_R are the polar angles counted from the centers of L and R Bose cloud, correspondingly (see Fig. 3). The angles $\phi_{L(R)}(\mathbf{r})$, defined as shown in Fig. 3, satisfy the conditions

$$\phi_R(\mathbf{r}_A) - \phi_L(\mathbf{r}_A) = \phi_L(\mathbf{r}_B) - \phi_R(\mathbf{r}_B) = \pi, \tag{46}$$

where $\mathbf{r}_A$ and $\mathbf{r}_B$ are the radius-vectors of Josephson links.

Substituting (44) into Eq. (43), using Eq. (46) and taking into account, that the functions Ψ_L and Ψ_R overlap in a small vicinity of A and B links, we obtain the following expression for the Josephson coupling parameter:

$$J = (J_A + J_B)\cos(\pi\Phi_{dr}) + i(J_A - J_B)\sin(\pi\Phi_{dr}), \tag{47}$$

where

$$J_{A(B)} \approx \int_{V_{A(B)}} d^3r\left[\frac{\hbar^2}{2m}\nabla|\Psi_L|\nabla|\Psi_R| + V_{tr}|\Psi_L||\Psi_R|\right]. \tag{48}$$



Figure 3. Left (L) and right(R) half-torus of the drag component, separated by Josephson links A and B.

Here V_A and V_B are the areas of overlapping of two one mode functions at links A and B, correspondingly.

Considering the Hilbert space in which the total number operator

$$\hat{N} = a^+_{\mathcal{L}} a_{\mathcal{L}} + a^+_{\mathcal{R}} a_{\mathcal{R}} \tag{49}$$

is a conservative quantity ($\hat{N} = N$) we present the Hamiltonian (40) in the following form

$$H_a = E_c(\hat{n}_{RL} + n_g)^2 + (J a^+_L a_R + h.c.) + const, \tag{50}$$

where

$$\hat{n}_{RL} = \frac{a^+_R a_R - a^+_L a_L}{2} \tag{51}$$

is the number difference operator,

$$E_c = \lambda_R + \lambda_L \tag{52}$$

is the interaction energy, and the quantity

$$n_g = \frac{1}{2E_c}\left[K_R - K_L + (N-1)(\lambda_R - \lambda_L)\right] \tag{53}$$

describes an asymmetry of L and R half-torus.

In what follows we imply that the system is in the Fock regime ($|J|N \ll E_c$) and use the number representation

$$|n_{RL}\rangle \equiv |n_R, n_L\rangle \equiv |\frac{N}{2} + n_{RL}, \frac{N}{2} - n_{RL}\rangle.$$

In this basis the first term in (50) is diagonal. The second term in (50) can be considered as small nondiagonal correction. But if n_g is biased near one of degeneracy points

$$n_{deg} = \begin{cases} M + \dfrac{1}{2} & \text{for even } N \\ M & \text{for odd } N \end{cases} \tag{54}$$

(where M is integer and $|M| < N/2$) the second term in (50) results in a strong mixing of two lowest states ($|\uparrow\rangle = |n_{deg} + 1/2\rangle$ and $|\downarrow\rangle = |n_{deg} - 1/2\rangle$) and the low energy dynamics of the system can be described by a pseudospin Hamiltonian

$$H_{eff} = -\frac{\Omega_x}{2}\hat{\sigma}_x - \frac{\Omega_y}{2}\hat{\sigma}_y - \frac{\Omega_z}{2}\hat{\sigma}_z, \tag{55}$$

where

$$\begin{aligned} \hat{\sigma}_x &= |\uparrow\rangle\langle\downarrow| + |\uparrow\rangle\langle\downarrow| \\ \hat{\sigma}_y &= -i(|\uparrow\rangle\langle\downarrow| - |\uparrow\rangle\langle\downarrow|) \\ \hat{\sigma}_z &= |\uparrow\rangle\langle\uparrow| - |\downarrow\rangle\langle\downarrow| \end{aligned} \tag{56}$$

are the Pauli operators, and

$$\begin{aligned} \Omega_x &= -(J_A + J_B)\sqrt{(N+1)^2 - n_{deg}^2}\cos(\pi\Phi_{dr}), \\ \Omega_y &= -(J_A - J_B)\sqrt{(N+1)^2 - n_{deg}^2}\sin(\pi\Phi_{dr}), \\ \Omega_z &= 2E_c(n_g - n_{deg}) \end{aligned} \tag{57}$$

are the components of the pseudomagnetic field.

If the quantities n_g, J_A and J_B depend on time and the conditions $N(J_A + J_B) \ll E_c$, $|n_g - n_{deg}| \ll 1$ are satisfied, our system can be used as qubit. In difference with the situation, considered in [3], our effective qubit Hamiltonian (55) contains the $\hat{\sigma}_y$ term. The appearance of this term in the Hamiltonian is caused by the drag effect. If the ratio J_A/J_B depends on time and the value of $2\Phi_{dr}$ is not integer this term cannot be excluded from the Hamiltonian by gauge transformation.

4. Detection of Berry phase in the Bose qubit

In experiments one can control the parameters n_g, J_A and J_B independently. It allows to realize an adiabatic cyclic evolution of the parameters of the Hamiltonian (55), for which vector $\mathbf{\Omega}$ subtends nonzero solid angle at the origin. Such an evolution results in appearance of a geometrical phase of the wave function known as Berry phase [19]. For the system (55) the value of this phase is the half of the solid angle subtending by the contour of evolution of $\mathbf{\Omega}$.

In this section we propose a scheme for generating Berry phase in the Bose qubit and show that in our scheme this phase is proportional to the value of Φ_{dr}, and, consequently, to the drag factor.

Let initially $J_A = J_B = 0$ and $n_g > n_{deg}$. The systems relaxes to the ground state $|\uparrow\rangle$. After that one switches on the equal Josephson coupling at links A and B ($J_A = J_B = J_{in}$) and sets the asymmetry parameter n_g equals to n_{deg}. The initial state of the qubit is the superposition of two instantaneous eigenstates $|e_a\rangle = (|\uparrow\rangle + |\downarrow\rangle)/\sqrt{2}$ and $|e_b\rangle = (|\uparrow\rangle - |\downarrow\rangle)/\sqrt{2}$:

$$|\psi_{in}\rangle = \frac{1}{\sqrt{2}}(|e_a\rangle + |e_b\rangle). \tag{58}$$

The expectation value of the relative number operator in the initial state $\langle\psi_{in}|\hat{n}_{RL}|\psi_{in}\rangle = 1/2$.

At the starting point the pseudomagnetic field is directed along the (-1,0,0) axes. Adiabatic evolution includes 4 stages.

At the 1-st stage the Josephson coupling J_B is switched off. After this stage pseudomagnetic field becomes aligned along the $(-\cos\pi\Phi_{dr}, -\sin\pi\Phi_{dr}, 0)$ direction.

At the 2-nd stage the coupling J_A is switched off and simultaneously the value of n_g is deviated from n_{deg}. After this stage $\mathbf{\Omega}$ is directed along the (0,0,1) axis.

At the 3-d stage the value of n_g is returned to the same degeneracy point and the coupling J_B is switched on. $\mathbf{\Omega}$ becomes directed in $(-\cos\pi\Phi_{dr}, \sin\pi\Phi_{dr}, 0)$ direction.

At the 4-th stage the coupling J_A is switched on and vector $\mathbf{\Omega}$ returns to its starting value.

After such an evolution the system arrives at the state

$$|\psi_m\rangle = \frac{1}{\sqrt{2}}\left(e^{i\delta_a + i\gamma}|e_a\rangle + e^{i\delta_b - i\gamma}|e_b\rangle\right), \tag{59}$$

where $\gamma = \pi\Phi_{dr}$ is the Berry phase equals to half of the solid angle subtending by $\mathbf{\Omega}$ and δ_a, δ_b are the dynamical phases.

Dynamical phases depends on time of evolution, while the Berry phase does not. Elimination of the dynamical phases is performed by swapping the eigenstates (π-transformation) and repeating the same cycle of evolution in a reverse direction [10].

The π-transformation can be done if one switches of the Josephson coupling and deviates the asymmetry parameter n_g from n_{deg} value during the time interval $t_\pi = 2\pi E_c(n_{deg} - n_g)/\hbar$. After the π-transformation the state of the qubit becomes

$$|\psi_{m\pi}\rangle = -\frac{i}{\sqrt{2}}(e^{i\delta_a + i\gamma}|e_b\rangle + e^{i\delta_b - i\gamma}|e_a\rangle). \tag{60}$$

After the cyclic evolution in the reverse direction the state becomes

$$|\psi_f\rangle = -\frac{i}{\sqrt{2}}e^{i(\delta_a+\delta_b)}(e^{2i\gamma}|e_b\rangle + e^{-2i\gamma}|e_a\rangle). \tag{61}$$

The expectation value of the difference number operator in the final state (61) differs from this value in the initial state (58) to the quantity that depends on the Berry phase and, consequently, on Φ_{dr}:

$$\langle\psi_f|\hat{n}_{RL}|\psi_f\rangle = \frac{1}{2} - \sin^2(2\gamma) = \frac{1}{2} - \sin^2(2\pi\Phi_{dr}). \tag{62}$$

Thus, the value of the drag can be determined from measurements of relative number of atoms in left and right condensates.

5. Conclusions

We derived a microscopic model of nondissipative drag in a two-component superfluid Bose gas and proposed possible implementation of a Bose qubit in which the drag effect is used to control quantum states. We have described how the Berry phase can be detected in such a qubit and found that this phase is proportional to the drag factor.

Main idea of our proposal is based on a correspondence between the drag force and a vector potential. One can found from Eq. (24) that the effective vector potential, applied to the drag component, is equal to $\mathbf{A}_{eff} = \hbar f_{dr}\nabla\varphi_{drv}$ (in units of $e = c = 1$), where φ_{drv} is the phase of the drive component, and f_{dr} is the drag factor, which is of order of the square root of the gas parameter. In a trap of a ring geometry the effective vector potential can be associated with an effective flux of external field $\Phi_{eff} = 2\pi\hbar f_{dr} N_v$, where N_v is the vorticity of the drive component. It allows to realize the Bose charge qubit, analogous to superconductive qubit, considered in [11]. Note that while Φ_{eff} is quantized, for $f_{dr} \ll 1$ it can be treated as a continuous one.

In conclusion we would mention another possibility that opens the drag effect. Tuning the effective flux Φ_{eff} to the $\Phi^{\pi}_{eff} = \pi\hbar$ value and increasing the Josephson coupling one can achieved a regime in which the states with different vorticities of the drag component are mixed. Such a system can be considered as a Bose counterpart of superconductive Josephson flux qubit.

Acknowledgements

This work is supported by the INTAS grant No 01-2344.

References

[1] Makhin Y, Schon G and Shnirman A 2001 *Rev. Mod. Phys.* **73** 357.
[2] Leggett A J 2001 *Rev. Mod. Phys.* **73** 307.
[3] Chen Z B and Zhang Y D 2002 *Phys. Rev.* A **65** 022318.
[4] Andreev A F and Bashkin E P 1975 *Zh. Exp. Teor. Fiz.* **69** 319 [Sov. Phys. JETP, **42** 164].
[5] Tanatar B and Das A K 1996 *Phys. Rev.* B **54** 13827.
[6] Terentjev S V and Shevchenko S I 1999 *Fiz. Nizk. Temp.* **25** 664 [*Low Temp. Phys.* **25** 493].
[7] Duan J M and Yip S 1993 *Phys. Rev. Lett.* **70** 3647.
[8] Duan J M 1995 *Europhysics Lett.* **29** 489.
[9] Rojo A G and Mahan G D 1992 *Phys. Rev. Lett.* **68** 2074.
[10] Ekert A, Ericsson M, Hayden P, Inamori H, Jones J A, Oi D K L and Vedral V 2000 *J. Mod. Opt.* **47** 2501.
[11] Falci G, Fazio R, Palma G M, Siewert J and Vedral V 2000 *Nature*(London) **407** 355.
[12] Shevchenko S I 1992 *Fiz. Nizk. Temp.* **18** 328 [*Sov. J. Low Temp. Phys.* **18** 223].
[13] Fil D V and Shevchenko S I 2001 *Phys. Rev.* A **64** 013607.
[14] Bogolyubov N N and Bogolyubov N N (Jr) 1982 *Introduction to quantum statistical mechanics* (Singapore: World Scientific).
[15] Stringari S 1996 *Phys. Rev. Lett.* **77** 2360.
[16] Öhberg P, Surkov E L, Tittonen I, Stenholm S, Wilkens M and Shlyapnikov G V 1997 *Phys. Rev.* A **56** R3346.
[17] Khawaja U Al, Pethick C J and Smith H 1999 *Phys. Rev.* A **60** 1507.
[18] Raghavan S, Smerzi A, Fantoni S and Shenoy S R 1999 *Phys. Rev.* A **59** 620.
[19] Berry M V 1984 *Proc. Roy. Soc.* A **392** 47.